

The intersection problem for small G -designs

ELIZABETH J. BILLINGTON*

*Centre for Combinatorics, Department of Mathematics,
The University of Queensland, Queensland 4072, AUSTRALIA*

DONALD L. KREHER

*Department of Mathematical Sciences, Michigan Technological University,
Houghton, Michigan, U.S.A. 49931-1295*

Abstract

A G -design of order n is a pair (P, B) where P is the vertex set of the complete graph K_n and B is an edge-disjoint decomposition of K_n into isomorphic copies of the simple graph G . Following design terminology, we call these copies "blocks". Given a particular graph G , the intersection problem asks for which k is it possible to find two G -designs (P, B_1) and (P, B_2) of order n , with $|B_1 \cap B_2| = k$, that is, with precisely k common blocks. Here we complete the solution of this intersection problem for several G -designs where G is "small", so that now it is solved for all connected graphs G with at most four vertices or at most four edges.


1 Introduction and preliminaries

Let G be a simple graph which is some subgraph of K_n , the complete undirected graph on n vertices. A G -design of order n is a pair (V, B) where V is the vertex set of K_n and B is an edge-disjoint decomposition of K_n into copies of the simple graph G . Following design terminology, we refer to these copies of G as blocks. Thus, for example, a Steiner triple system is a K_3 -design and a balanced incomplete block design with block size four and index $\lambda = 1$ is a K_4 -design. The number of blocks, $|B|$, is $\binom{n}{2}/|E(G)|$ where $E(G)$ is the edge-set of G ; this number clearly must be an integer.

*Research supported by the Australian Research Council.

The intersection problem for G -designs asks for two designs based on the same set V , with $|B_1 \cap B_2| = k$; that is, having precisely k common blocks. This problem was first considered for Steiner triple systems or K_3 -designs (see [8]), and since then the intersection problem has been considered for many different types of combinatorial structures; see [3] for a recent survey.




A (p, q) graph is one with p vertices and q edges. We list below all non-trivial connected simple (p, q) graphs with $\min(p, q) \leq 4$.




Clearly a K_2 -design is unique; each block is an edge! And so for this design we cannot find two distinct designs, let alone a pair of designs intersecting in a specified number of blocks! So we leave this trivial case.

As mentioned above, the intersection problem for K_3 -designs was dealt with in [8]. The intersection problem for C_4 -designs appears in [4], for $(K_4 - e)$ -designs in [5] and for K_4 -designs (with a few exceptions) in [6].

The remaining cases, namely the graphs $P_3, P_4, P_5, S_3, S_4, D$ and Y , we deal with below. We use the notation of [2] for names of these graphs, and the following diagram shows how we label the blocks.





In what follows we let $IG(n)$ denote the set of integers k for which there exist two G -designs (P, B_1) and (P, B_2) with $|P| = n$ and $|B_1 \cap B_2| = k$. Also if G is a graph with q edges, let

$$JG(n) = \begin{cases} \{0, 1, 2, \dots, \frac{1}{q} \binom{n}{2} - 2, \frac{1}{q} \binom{n}{2}\} & \text{if } q | \binom{n}{2}; \\ \emptyset & \text{otherwise.} \end{cases}$$

In other words, $JG(n)$ denotes the intersection numbers one expects to achieve with a G -design of order n .

We also modify this notation slightly and let $IG(H)$ and $JG(H)$ denote respectively the achievable and expected intersection numbers for a G -decomposition of the graph H .

We also need the following definition. If S is a set of positive integers and h is some positive integer, then $h * S$ denotes the set of all integers which can be obtained by adding any h elements of S together (repetitions of elements of S allowed). For example, $2 * \{0, 1, 3\} = \{0, 1, 2, 3, 4, 6\}$.

Subsequently we shall need to decompose certain bipartite and tripartite graphs into edge-disjoint copies of the graphs G . Consider the following example.

EXAMPLE 1.1 *Decompositions of $K_{4,4}$ into copies of P_5 .*

Let $K_{4,4}$ have vertex set $\{1_1, 2_1, 3_1, 4_1\} \cup \{1_2, 2_2, 3_2, 4_2\}$, and let $P = \{A, B, C, D\}$ where

$$\begin{aligned} A &= (1_2, 1_1, 2_2, 2_1, 3_2), & B &= (1_2, 4_1, 4_2, 3_1, 3_2), \\ C &= (1_1, 3_2, 4_1, 2_2, 3_1), & D &= (1_1, 4_2, 2_1, 1_2, 3_1). \end{aligned}$$

These cover the 16 edges of $K_{4,4}$, and so form a P_5 -decomposition of $K_{4,4}$.

Now C and D cover the same edges as

$$C' = (1_2, 2_1, 4_2, 1_1, 3_2), \quad D' = (1_2, 3_1, 2_2, 4_1, 3_2),$$

while B, C and D together cover the same edges as

$$\hat{B} = (2_2, 4_1, 3_2, 1_1, 4_2), \quad \hat{C} = (1_2, 2_1, 4_2, 3_1, 3_2), \quad \hat{D} = (2_2, 3_1, 1_2, 4_1, 4_2).$$

Moreover, the permutation (1 2) applied to the subscripts of blocks A, B, C and D yields a different P_5 -decomposition of $K_{4,4}$ having no blocks in common with P ; call these blocks \bar{P} .

Thus we see that $|P \cap \bar{P}| = 0$, $|P \cap \{A, \hat{B}, \hat{C}, \hat{D}\}| = 1$, $|P \cap \{A, B, C', D'\}| = 2$, $|P \cap P| = 4$. (Clearly it is not possible to have two decompositions which have all but one block in common.) We record these intersection numbers for P_5 -decompositions of $K_{4,4}$ as

$$IP_5(K_{4,4}) = \{0, 1, 2, 4\}. \quad \square$$

More generally, if \mathcal{K} is a collection of graphs, then a \mathcal{K} -decomposition of the graph H , (V, \mathcal{B}) , is an edge-disjoint decomposition of H with vertex set V into a set of subgraphs \mathcal{B} , with each subgraph isomorphic to some graph in \mathcal{K} . If $\mathcal{K} = \{G\}$, then we call this a G -decomposition of H , and if also $H = K_n$, then it is a G -design of order n .

The following lemma will be most useful in the rest of this paper.

LEMMA 1.1 *Let G be a graph with q edges and suppose (V, \mathcal{B}) is a $\{K_m, H\}$ -decomposition of K_n , with $\alpha > 0$ blocks isomorphic to K_m . If $IG(m) = JG(m)$ and $IG(H) \supseteq \{0, r\}$ with $|E(H)| = qr$ and $q(r+1) \leq \alpha \binom{m}{2}$, then $IG(n) = JG(n)$.*

Proof. First a G -design of order n can be constructed by replacing each of the blocks $B \in \mathcal{B}$ that is isomorphic to K_m by a G -design of order m , and replacing each of the blocks $B \in \mathcal{B}$ that is isomorphic to H by a G -decomposition of H .

Secondly, if $q \mid \binom{m}{2}$, then for any positive integer x ,

$$x * JG(m) = \left\{ 0, 1, 2, \dots, \frac{x}{q} \binom{m}{2} - 2, \frac{x}{q} \binom{m}{2} \right\},$$

and for all $x \geq r+1$,

$$\{0, 1, 2, \dots, x-2, x\} + \{0, r\} = \{0, 1, 2, \dots, x+r-2, x+r\}.$$

Thus if \mathcal{B} contains α blocks isomorphic to K_m and β blocks isomorphic to H , then

$$IG(v) \supseteq \alpha * JG(m) + \beta * \{0, r\} = \{0, 1, 2, \dots, z-2, z\}$$

where $z = \alpha \frac{1}{q} \binom{m}{2} + \beta r$. But \mathcal{B} is a decomposition of K_n so we also have $\alpha \binom{m}{2} + \beta qr = \binom{n}{2}$. Thus $z = \frac{1}{q} \binom{n}{2}$, as required. Hence $IG(n) = JG(n)$. \square

In what follows, the graph H in Lemma 1.1 will usually be a complete bipartite or tripartite graph.

2 Paths on 3, 4 and 5 vertices

2.1 The path P_3

Note that a P_3 -design of order n contains $n(n-1)/4$ blocks and so we must have $n \equiv 0$ or $1 \pmod{4}$.

EXAMPLE 2.1 $IP_3(K_{2,2}) = \{0, 2\}$.

Take designs (P, B_i) , $i = 1, 2$, where the vertex set of $K_{2,2}$ is $P = \{a, b\} \cup \{c, d\}$, and $B_1 = \{(a, c, b), (a, d, b)\}$, $B_2 = \{c, a, d\}, (c, b, d)\}$. Since $|B_1 \cap B_2| = 0$ we have $IP_3(K_{2,2}) = \{0, 2\}$. \square

EXAMPLE 2.2 $IP_3(4) = \{0, 1, 3\}$.

We use designs (P, B_i) , $i = 1, 2, 3$, where $P = \{a, b, c, d\}$ and

$$\begin{aligned} B_1 &= \{(a, b, c), (a, c, d), (a, d, b)\}, \\ B_2 &= \{(a, b, c), (d, a, c), (b, d, c)\}, \\ B_3 &= \{(a, b, d), (a, d, c), (a, c, b)\}. \end{aligned}$$

Here $|B_1 \cap B_2| = 1$, $|B_1 \cap B_3| = 0$ and of course $|B_1 \cap B_1| = 3$. The result follows. \square

EXAMPLE 2.3 $IP_3(K_{1,2n}) = \{0, 1, 2, \dots, n-2, n\}$.

The verification of this is immediate. \square

Now let $n = 4m$, and take the vertex set of K_n to be $\{(i, j) \mid 1 \leq i \leq 2m, j = 1, 2\}$. Take K_4 blocks $\{(2i-1, j), (2i, j) \mid j = 1, 2\}$, for $1 \leq i \leq m$, and $K_{2,2}$ blocks $\{(a, 1), (a, 2)\} \cup \{(b, 1), (b, 2)\}$ where $1 \leq a < b \leq 2m$ and $\{a, b\} \neq \{2i-1, 2i\}$ for $1 \leq i \leq m$. The result is a $\{K_4, K_{2,2}\}$ -decomposition of K_{4m} and consequently by Lemma 1.1 we have $IP_3(4m) = JP_3(4m)$.

Now let $n = 4m + 1$, and let the vertex set of K_n be $\{1, 2, \dots, 4m, \infty\}$. We may use P_3 -designs of order $4m$ on $\{1, 2, \dots, 4m\}$ and use Example 2.3 to find P_3 -decompositions of $K_{1,4m}$ on $\{\infty\} \cup \{1, 2, \dots, 4m\}$. Thus

$$\begin{aligned} IP_3(4m+1) &\supseteq IP_3(4m) + IP_3(K_{1,4m}) \\ &= \{0, 1, 2, \dots, m(4m+1) - 2, m(4m+1)\} = JP_3(4m+1). \end{aligned}$$

We have now proved

THEOREM 2.1 *The intersection numbers for P_3 -designs are given by $IP_3(n) = JP_3(n) = \{0, 1, \dots, b-2, b\}$ where $b = n(n-1)/4$, the total number of blocks in a P_3 -design of order n .* \square

2.2 The path P_4

A P_4 -design of order n contains $n(n-1)/6$ blocks so that $n \equiv 0$ or $1 \pmod{3}$, $n \geq 4$. So let $n = 3m$ or $3m + 1$. First we give some necessary examples.

EXAMPLE 2.4 $IP_4(4) = \{0, 2\}$.

Let $V = \{1, 2, 3, 4\}$, $B_1 = \{(1, 2, 3, 4), (2, 4, 1, 3)\}$, $B_2 = \{(1, 4, 3, 2), (3, 1, 2, 4)\}$. Then (V, B_1) , (V, B_2) are both P_4 -designs, and $|B_1 \cap B_2| = 0$; the result follows. \square

EXAMPLE 2.5 $IP_4(K_{3,3}) \supseteq \{0, 3\}$.

Let $K_{3,3}$ have vertex set $V = \{1, 2, 3\} \cup \{4, 5, 6\}$. Two disjoint decompositions are $B_1 = \{(1, 4, 2, 5), (2, 6, 3, 4), (3, 5, 1, 6)\}$, $B_2 = \{(2, 5, 3, 6), (3, 4, 1, 5), (1, 6, 2, 4)\}$. The result follows. \square

EXAMPLE 2.6 $IP_4(6) = \{0, 1, 2, 3, 5\}$.

Let K_6 have vertex set $V = \{0, 1, 2, 3, 4, 5\}$, and let $A = \{(0, 1, 2, 3), (3, 0, 5, 2), (0, 4, 3, 1)\}$, $B = \{(0, 2, 4, 5), (3, 5, 1, 4)\}$, and $C = \{(0, 4, 3, 1), (3, 5, 1, 4)\}$. Then $(V, A \cup B)$ is one P_4 -design of order 6. Note that the blocks A trade with $A' = \{(1, 0, 5, 2), (4, 0, 3, 2), (4, 3, 1, 2)\}$, and the blocks C trade with $C' = \{(0, 4, 1, 3), (1, 5, 3, 4)\}$. Let $X = A \cup B$, and let α denote the permutation $(14)(35)$ and β the permutation $(15)(34)$. The following table lists intersection numbers.

blocks	intersection size
$X, X\alpha$	0
$X, X\beta$	1
$X, \{A', B\}$	2
$X, ((X \setminus C) \cup C')$	3
X, X	5

□

EXAMPLE 2.7 $IP_4(7) = \{0, 1, 2, 3, 4, 5, 7\}$.

Let K_7 have vertex set $V = \{0, 1, 2, 3, 4, 5, 6\}$. Let $A = \{(0, 1, 3, 6), (1, 2, 4, 0)\}$, $B = \{(2, 3, 5, 1), (3, 4, 6, 2), (4, 5, 0, 3)\}$ and $C = \{(5, 6, 1, 4), (6, 0, 2, 5)\}$. Then (V, X) , where $X = (A \cup B \cup C)$, is a P_4 -design of order 7. Moreover, A, B and C trade with $A' = \{(2, 1, 3, 6), (1, 0, 4, 2)\}$, $B' = \{(1, 5, 0, 3), (6, 4, 5, 3), (6, 2, 3, 4)\}$ and $C' = \{(4, 1, 6, 0), (0, 2, 5, 6)\}$ respectively. Let α denote the permutation $(06)(13)$. The following table lists intersection numbers.

blocks	intersection size
$X, A' \cup B' \cup C'$	0
$X, X\alpha$	1
$X, A \cup B' \cup C'$	2
$X, A' \cup B \cup C'$	3
$X, A \cup B' \cup C$	4
$X, A' \cup B \cup C$	5
X, X	7

□

EXAMPLE 2.8 $IP_4(9) = \{0, 1, 2, \dots, 9, 10, 12\}$.

Take a P_4 -design of order 6, on $\{0, 1, 2, 3, 4, 5\}$, and adjoin elements H, J and K , and also the blocks

$$X = \{(0, H, 1, J), (2, H, 3, J), (4, H, 5, J), (1, K, 0, J), \\ (3, K, 2, J), (5, K, J, H), (H, K, 4, J)\}.$$

Now using $IP_4(6)$ we have $\{7, 8, 9, 10, 12\} \subseteq IP_4(9)$. Also applying the permutation (HJ) to the set X changes all the blocks in X , so again using $IP_4(6)$ we have $\{0, 1, 2, 3, 5\} \subseteq IP_4(9)$.

Thus it remains to show that 4 and 6 are as in $IP_4(9)$. To do this, let D denote the design with blocks $X \cup A \cup B$ where A and B are as in Example 2.6 above. Then $|D \cap D\gamma| = 4$ where γ is the permutation $(03)(12)$. Finally, let

$$T = \{(0, 2, 4, 5), (3, 0, 5, 2), (0, 4, 3, 1), (3, 5, 1, 4)\}$$

which has trade

$$T' = \{(3, 1, 4, 5), (0, 3, 5, 1), (3, 4, 2, 0), (4, 0, 5, 2)\}.$$

Then $|D\gamma \cap ((D \setminus T) \cup T')| = 6$, which completes the intersection numbers for designs of order 9. \square

Now let $n = 3m + 1$ and let the vertex set of K_n be $V = \{(i, j) \mid 1 \leq i \leq m, j = 1, 2, 3\} \cup \{\infty\}$. There is a $\{K_7, K_4, K_{3,3}\}$ -decomposition of K_n with: one K_7 block $\{\infty\} \cup \{(i, j) \mid i = 1, 2; j = 1, 2, 3\}$; K_4 blocks $\{\infty\} \cup \{(i, j) \mid j = 1, 2, 3\}$, for $3 \leq i \leq m$; $K_{3,3}$ blocks $\{(i, j) \mid j = 1, 2, 3\} \cup \{(i', j) \mid j = 1, 2, 3\}$, for all $1 \leq i < i' \leq m$, excluding $\{i, i'\} = \{1, 2\}$. Then using Examples 2.7, 2.4, 2.5 and a slight generalization of Lemma 1.1, it follows that $IP_4(3m + 1) = JP_4(3m + 1) = \{0, 1, 2, \dots, t - 2, t\}$ where $t = m(3m + 1)/2$, the total number of blocks in a P_4 -design of order $3m + 1$.

Next let $n = 3m$. The cases m even and m odd are treated separately. When m is even let $n = 6M$ and let the vertex set of K_n be $\{(i, j) \mid 1 \leq i \leq 2M; j = 1, 2, 3\}$. There is a $\{K_6, K_{3,3}\}$ -decomposition of K_n with K_6 blocks $\{(2i - 1, j), (2i, j) \mid j = 1, 2, 3\}$ for $1 \leq i \leq M$ and $K_{3,3}$ blocks $\{(i_1, j) \mid j = 1, 2, 3\} \cup \{(i_2, j) \mid j = 1, 2, 3\}$ for all $1 \leq i_1 < i_2 \leq 2M$ excluding $\{i_1, i_2\} = \{2i - 1, 2i\}$, $1 \leq i \leq M$.

The result $IP_4(6M) = JP_4(6M)$ then follows from Examples 2.6, 2.5 and Lemma 1.1.

When m is odd let $n = 6M + 3$, and let the vertex set of K_n be $\{(i, j) \mid 1 \leq i \leq 2M + 1, j = 1, 2, 3\}$. There is a $\{K_9, K_6, K_{3,3}\}$ -decomposition of K_n with: one K_9 block $\{(i, j) \mid i, j = 1, 2, 3\}$; K_6 blocks $\{(2i, j), (2i + 1, j) \mid j = 1, 2, 3\}$ for $i = 2, \dots, M$; $K_{3,3}$ blocks $\{(a, j) \mid j = 1, 2, 3\} \cup \{(b, j) \mid j = 1, 2, 3\}$ for all pairs $\{a, b\}$ with $a \neq b$ and with a and b not both in $\{1, 2, 3\}$ or in $\{2i, 2i + 1\}$, $2 \leq i \leq M$.

Then from Examples 2.8, 2.6, 2.5 and Lemma 1.1, we have $IP_4(6M + 3) = JP_4(6M + 3)$.

We have now proved

THEOREM 2.2 *The intersection numbers for P_4 -designs are given by $IP_4(n) = \{0, 1, \dots, b - 2, b\}$ where $b = n(n - 1)/6$. \square*

2.3 The path P_5

The graph P_5 has 4 edges, and so a suitable decomposition of K_n will contain $n(n - 1)/8$ blocks; consequently we must have $n \equiv 0$ or $1 \pmod{8}$. The only ingredients needed are decompositions of $K_{4,4}$, K_8 and K_9 , and of course their intersection numbers too.

Now let the vertex set of K_n be $V = \{(i, j) \mid 1 \leq i \leq 2m, 1 \leq j \leq 4\} \cup V \cup \{\infty\}$, according as $n = 8m$ or $8m + 1$.

In the former case there is a $\{K_8, K_{4,4}\}$ -decomposition of K_n with K_8 blocks $\{(2i - 1, j), (2i, j) \mid 1 \leq j \leq 4\}$ for $1 \leq i \leq m$, and $K_{4,4}$ blocks $\{(a, j) \mid 1 \leq j \leq 4\} \cup \{(b, j) \mid 1 \leq j \leq 4\}$ for all $1 \leq a < b \leq 2m$ and $\{a, b\} \neq \{2i - 1, 2i\}$ for $1 \leq i \leq m$. In the latter case there is a $\{K_9, K_{4,4}\}$ -decomposition of K_n ; the K_9 blocks have $\{\infty\}$ adjoined to each of the K_8 blocks above, otherwise blocks are the same as when $n = 8m$.

In Example 1.1 we showed that $IP_5(K_{4,4}) = \{0, 1, 2, 4\}$. We also need the following two examples.

EXAMPLE 2.9 $IP_5(8) = \{0, 1, 2, 3, 4, 5, 7\}$.

On the vertex set $\mathbb{Z}_7 \cup \{\infty\}$, developing the base block $\beta = (\infty, 0, 1, 3, 6)$ modulo 7 generates a P_5 -decomposition of K_7 . For each $i \in \mathbb{Z}_7$ the blocks $A_i = \{\beta + i, \beta + i + 1\}$ trade with $A'_i = \{(6, 3, 1, 0, 4) + i, (0, \infty, 1, 2, 4) + i\}$, and $B = \{\beta + 4, \beta + 5, \beta + 6\}$ trades with $B' = \{(3, 0, 5, \infty, 4), (\infty, 6, 5, 2, 0), (0, 6, 1, 4, 5)\}$. We observe that A_0, A_2 and A_4 are mutually disjoint and that B is disjoint from A_0 and A_2 . Consequently $IP_5(8) = \{0, 1, 2, 3, 4, 5, 7\}$. \square

EXAMPLE 2.10 $IP_5(9) = \{0, 1, 2, 3, 4, 5, 6, 7, 9\}$.

On the vertex set \mathbb{Z}_9 , a P_5 -design is generated by developing the base block $\beta = (0, 1, 3, 7, 4)$ (modulo 9). For each $i \in \mathbb{Z}_9$ the blocks $A_i = \{\beta + i, \beta + i + 2\}$ trade with $A'_i = \{(0, 5, 3, 7, 4) + i, (2, 3, 1, 0, 6) + i\}$ and the blocks $B_i = \{\beta + i, \beta + i + 1, \beta + i + 2\}$ trade with $B'_i = \{(0, 1, 2, 3, 5) + i, (1, 3, 7, 4, 2) + i, (6, 0, 5, 8, 4) + i\}$.

Moreover, the blocks $C = \{\beta + 5, \beta + 7, \beta + 8\}$ trade with the blocks $C' = \{(7, 8, 3, 6, 5), (3, 0, 8, 6, 2), (8, 1, 5, 2, 0)\}$. The following table lists the disjoint trades which may be used in order to achieve the required intersection values.

trades	intersection achieved
B_0, B_3, B_6	0
A_0, A_1, A_4, A_5	1
C, A_0, A_1	2
B_0, B_3	3
A_0, B_6	4
A_0, A_1	5
B_0	6
A_0	7
nothing	9

\square

Now applying Lemma 1.1 yields the following result for P_5 -designs.

THEOREM 2.3 *The intersection numbers for P_5 -designs are given by $IP_5(n) = \{0, 1, \dots, b - 2, b\}$ where $b = n(n - 1)/8$.* \square

3 Stars with 3 and 4 edges

3.1 S_3 -designs

The number of blocks in an S_3 -design of order n is $n(n-1)/6$, and so $n \equiv 0$ or $1 \pmod{3}$, and $n \geq 6$. (S_3 involves four vertices, and it is easy to see that K_4 has no S_3 -decomposition.)

We start with the following example.

EXAMPLE 3.1 $IS_3(K_{3,3}) = \{0, 3\}$.

Let $K_{3,3}$ have vertex set $\{1, 2, 3\} \cup \{4, 5, 6\}$. The following two S_3 -decompositions are disjoint.

$$D_1 = \{(1 : 4, 5, 6), (2 : 4, 5, 6), (3 : 4, 5, 6)\},$$

$$D_2 = \{(4 : 1, 2, 3), (5 : 1, 2, 3), (6 : 1, 2, 3)\}.$$

Moreover, it is straightforward to see that $1 \notin IS_3(K_{3,3})$. □

One slight difficulty in this case (and, indeed, for S_m -designs in general) is that the expected full set of intersection numbers for a design of order 6 (or $2m$ in general) cannot be achieved. In the case of S_3 -designs, each block involves 4 vertices, and it is impossible to find a trade consisting of two blocks when the design is of order 6. The smallest trade involves *seven* vertices, such as $\{(x : a, b, c), (x : d, e, f)\}$ trading with $\{(x : a, b, d), (x : c, e, f)\}$. We do however achieve the other expected intersection numbers, as the following example shows.

EXAMPLE 3.2 $IS_3(6) = \{0, 1, 2, 5\}$.

Let $V = \{0, 1, 2, 3, 4, 5\}$ and take

$$B = \{(0 : 5, 1, 2), (1 : 5, 2, 3), (2 : 5, 3, 4), (3 : 5, 4, 0), (4 : 5, 0, 1)\}.$$

Let $\alpha = (012)$, $\beta = (345)$ and $\gamma = (01)$ be permutations on V . The result then follows from the table below.

blocks	intersection
$B \cap B\alpha$	0
$B \cap B\beta$	1
$B \cap B\gamma$	2
$B \cap B$	5

□

Three more necessary examples follow.

EXAMPLE 3.3 $IS_3(7) = \{0, 1, 2, 3, 4, 5, 7\}$.

Take the vertex set $\{0, 1, 2, 3, 4, 5, 6\}$, and blocks $B \cup \{(6 : 0, 1, 2), (6 : 3, 4, 5)\} = B \cup Y$ where B is as in Example 3.2. The permutations α, β and γ of Example 3.2 fix Y . Hence $\{2, 3, 4, 7\} \subseteq IS_3(7)$. Moreover, Y trades with $Y' = \{(6 : 0, 1, 3), (6 : 2, 4, 5)\}$, and so $0 \in IS_3(7)$. Also $|(B \cup Y) \cap (B\beta \cup Y')| = 1$ and $|(B \cup Y) \cap (B \cup Y')| = 5$, so the result follows. \square

EXAMPLE 3.4 $IS_3(9) = \{0, 1, \dots, 10, 12\}$.

Let the vertex set be \mathbb{Z}_9 , and take blocks B as follows.

block	in subset(s)	block	in subset(s)
$(0 : 1, 3, 6)$	X	$(6 : 1, 2, 7)$	Y, T
$(1 : 2, 4, 7)$	Y	$(7 : 2, 0, 8)$	T
$(2 : 0, 5, 8)$		$(8 : 0, 1, 6)$	X, T
$(3 : 1, 4, 6)$	X	$(3 : 2, 7, 8)$	
$(4 : 2, 5, 7)$	Y	$(4 : 0, 8, 6)$	X
$(5 : 0, 3, 8)$	Z	$(5 : 1, 6, 7)$	Y, Z

The set X trades with $X' = \{(1 : 0, 3, 8), (0 : 3, 4, 8), (6 : 0, 3, 8), (4 : 3, 6, 8)\}$; the set Y trades with $Y' = \{(2 : 1, 4, 6), (1 : 4, 5, 6), (7 : 1, 4, 6), (5 : 4, 6, 7)\}$; the set Z trades with $Z' = \{(5 : 0, 3, 7), (5 : 8, 1, 6)\}$; and the set T trades with $T' = \{(6 : 1, 2, 8), (7 : 0, 2, 6), (8 : 0, 1, 7)\}$. Also Z and T are disjoint. The intersection values now follow from the table below, where numbers in parentheses are permutations on \mathbb{Z}_9 .

blocks	intersection
$B \cap B(678)$	0
$B \cap B(4758)$	1
$B \cap B(45)(78)$	2
$B \cap B(78)$	3
$B \cap ((B \setminus (X \cup Y)) \cup X' \cup Y')$	4
$B \cap B(45)$	5
$B \cap ((B \setminus (X \cup Z)) \cup X' \cup Z')$	6
$B \cap ((B \setminus (X \cup T)) \cup X' \cup T')$	7
$B \cap ((B \setminus X) \cup X')$	8
$B \cap ((B \setminus T) \cup T')$	9
$B \cap ((B \setminus Z) \cup Z')$	10
$B \cap B$	12

\square

EXAMPLE 3.5 $IS_3(10) = \{0, 1, \dots, 13, 15\}$.

Take \mathbb{Z}_{10} and blocks B of Example 3.4 above, together with $P = \{(9 : 0, 1, 2), (9 : 3, 4, 5), (9 : 6, 7, 8)\}$. The blocks in P are fixed by the above permutations (except for (4758)) and by the trades on B , so $\{3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 15\} \subseteq IS_3(10)$. Also P trades with $P' = \{(9 : 0, 1, 3), (9 : 2, 4, 6), (9 : 5, 7, 8)\}$, and so in particular $\{0, 1, 5\} \subseteq IS_3(10)$ also. Finally we see that $2 \in IS_3(10)$, using $1 \in IS_3(9)$ and the

trade $\{(9 : 0, 1, 2), (9 : 3, 4, 5)\}$ with $\{(9 : 0, 1, 3), (9 : 2, 4, 5)\}$. This completes the example. \square

In the general situation we deal with four cases: $n = 6m$, $n = 6m + 1$, $n = 6m + 3$ and $n = 6m + 4$. In each case the vertex set is $V = \{(i, j) \mid 1 \leq i \leq 2m, j = 1, 2, 3\}$, or $V \cup \{\infty\}$, or $V' = V \cup \{(2m + 1, j) \mid j = 1, 2, 3\}$ or $V' \cup \{\infty\}$ (respectively).

First, when $n = 6m + 1$, there is a $\{K_7, K_{3,3}\}$ -decomposition of K_n with K_7 blocks $\{\infty\} \cup \{(2i - 1, j), (2i, j) \mid j = 1, 2, 3\}$ for $1 \leq i \leq m$ and $K_{3,3}$ blocks $\{(a, j) \mid j = 1, 2, 3\} \cup \{(b, j) \mid j = 1, 2, 3\}$, for all $1 \leq a < b \leq 2m$, excluding $\{a, b\} = \{2i - 1, 2i\}, 1 \leq i \leq m$.

From Lemma 1.1 it follows that $IS_3(6m + 1) = JS_3(6m + 1)$.

Secondly, when $n = 6m + 4$, we use a $\{K_{10}, K_7, K_{3,3}\}$ -decomposition of K_n , with one K_{10} block and $m - 1$ K_7 blocks. Once again Lemma 1.1 then yields $IS_3(6m + 4) = JS_3(6m + 4)$.

Thirdly, when $n = 6m$, in order to achieve the intersection number " $b - 2$ ", with all but two blocks in common, since $5 - 2 = 3 \notin IS_3(6)$, we use a $\{K_9, K_6, K_{3,3}\}$ -decomposition of K_n with two K_9 blocks and $m - 3$ K_6 blocks. This assumes that $m \geq 3$, so $n \geq 18$; the case of order 12, therefore, must be considered separately.

Then, for $m \geq 3$, as before we obtain $IS_3(6m) = JS_3(6m)$.

Fourthly, when $n = 6m + 3$, we use a $\{K_9, K_6, K_{3,3}\}$ -decomposition of K_n with one K_9 block and $m - 1$ K_6 blocks, and obtain $IS_3(6m + 3) = JS_3(6m + 3)$.

It now remains to consider the case of order 12.

EXAMPLE 3.6 $IS_3(12) = \{0, 1, \dots, 20, 22\}$.

First, all intersection numbers except 20 (that is, $(b - 2)$) can be achieved with the following construction using two designs of order 6 and four lots of decompositions of $K_{3,3}$. Let A, B, C and D each stand for a set of three vertices. Then on sets $\{A, B\}$ and $\{C, D\}$, place S_3 -designs of order 6, and on the sets $\{A\} \cup \{C\}$, $\{A\} \cup \{D\}$, $\{B\} \cup \{C\}$, and $\{B\} \cup \{D\}$, place S_3 -decompositions of $K_{3,3}$. The result is an S_3 -design of order 12, and we see that

$$IS_3(12) \supseteq 2 * IS_3(6) + 4 * IS_3(K_{3,3})$$

which includes all required intersection numbers except 20.

Secondly, in order to obtain this intersection number, note that in the above construction, one of the four decompositions of $K_{3,3}$ is on the sets $\{A\} \cup \{C\}$ while another is on the sets $\{A\} \cup \{D\}$; so there will be two blocks of the form $(x : u, v, w)$ and $(x : r, s, t)$. These may be traded with $(x : u, v, t)$ and $(x : r, s, w)$; so we have $20 \in IS_3(12)$ as required. \square

The results in this subsection have shown

THEOREM 3.1 *The intersection numbers for S_3 -designs are given by $IS_3(n) = \{0, 1, \dots, b - 2, b\}$ where $n \equiv 0$ or $1 \pmod{3}$, $n \geq 6$ and $b = n(n - 1)/6$, except that $3 \notin IS_3(6)$.* \square

3.2 S_4 -designs

Since the number of blocks in an S_4 -design of order n is $n(n-1)/8$, we must have $n \equiv 0$ or $1 \pmod{8}$. First note that once we have intersection numbers $IS_4(8m)$, we can easily obtain $IS_4(8m+1)$. For in order to construct an S_4 -design of order $8m+1$ from one of order $8m$ we may simply adjoin one new vertex, say x , and $2m$ new blocks of the form $\{(x : a, b, c, d) \mid a, b, c, d \in V\}$ where V is the vertex set of the design of order $8m$. Moreover, by judicious interchange of the $2m$ elements, we see that we may construct two S_4 -designs of order $8m+1$ so that

$$IS_4(8m+1) \supseteq IS_4(8m) + \{0, 1, 2, \dots, 2m-2, 2m\}.$$

Now consider the following examples.

EXAMPLE 3.7 $IS_4(K_{4,4}) \supseteq \{0, 4\}$.

Imitate the construction in Example 3.1 above, but taking four vertices rather than three in each partite set. □

EXAMPLE 3.8 $IS_4(8) = \{0, 1, 2, 3, 4, 7\}$.

With vertex set $\{0, 1, 2, 3, 4, 5, 6, 7\}$, let blocks B be as follows.

$$\begin{aligned} &(0 : 1, 2, 3, 7), (1 : 2, 3, 4, 7), (2 : 3, 4, 5, 7), (3 : 4, 5, 6, 7), \\ &(4 : 5, 6, 0, 7), (5 : 6, 0, 1, 7), (6 : 0, 1, 2, 7). \end{aligned}$$

The following table shows the intersection values achieved by applying the given permutations to the vertices.

permutation	intersection size
(0 1 2 3)	0
(∞ 0 1 2)	1
(0 1 2)	2
(∞ 0)	3
(0 1)	4
identity	7

□

EXAMPLE 3.9 $IS_4(9) = \{0, 1, 2, 3, 4, 5, 6, 7, 9\}$.

As indicated in the remark preceding Example 3.7,

$$\begin{aligned} IS_4(9) &\supseteq IS_4(8) + \{0, 2\} \\ &= \{0, 1, 2, 3, 4, 7\} + \{0, 2\} \\ &= \{0, 1, 2, 3, 4, 5, 6, 7, 9\}. \end{aligned}$$

□

First, all intersection numbers except 28 (that is, $(b - 2)$) can be achieved with the following construction using two designs of order 8 and four lots of decompositions of $K_{4,4}$. Let A, B, C and D each stand for a set of four vertices. Then on sets $\{A, B\}$ and $\{C, D\}$, place S_4 -designs of order 8, and on the sets $\{A\} \cup \{C\}$, $\{A\} \cup \{D\}$, $\{B\} \cup \{C\}$, and $\{B\} \cup \{D\}$, place S_4 -decompositions of $K_{4,4}$. The result is an S_4 -design of order 16, and we see that

$$IS_4(16) \supseteq 2 * IS_4(8) + 4 * IS_4(K_{4,4})$$

which includes all required intersection numbers except 28.

Secondly, in order to obtain this intersection number, take another design of order 16 with vertex set $\mathbb{Z}_{15} \cup \{\infty\}$ and 30 blocks as follows:

$$(i : i + 1, i + 2, i + 3, i + 4), (i : i + 5, i + 6, i + 7, \infty), \quad i \in \mathbb{Z}_{15}.$$

The two blocks $(0 : 1, 2, 3, 4)$, $(0 : 5, 6, 7, \infty)$ trade with $(0 : 5, 6, 7, 4)$, $(0 : 1, 2, 3, \infty)$, changing just two blocks, and thus showing that $28 \in IS_4(16)$ as required. \square

Again, using the remark at the start of this subsection, using the above example it is easy to obtain $IS_4(17) = \{0, 1, \dots, 32, 34\}$.

Now the general construction for order $8m$ uses a $\{K_{16}, K_8, K_{4,4}\}$ -decomposition of K_{8m} with one K_{16} block and $m - 2$ K_8 blocks. Explicitly, let the vertex set be $\{(i, j) \mid 1 \leq i \leq 2m, 1 \leq j \leq 4\}$, and let the K_{16} block be $\{(i, j) \mid 1 \leq i, j \leq 4\}$, the K_8 blocks be $\{(2i - 1, j), (2i, j) \mid 1 \leq j \leq 4\}$ for $3 \leq i \leq m$, and the $K_{4,4}$ blocks be $\{(a, j) \mid 1 \leq j \leq 4\} \cup \{(b, j) \mid 1 \leq j \leq 4\}$ for all $a \neq b$ where a and b are not both first components of elements in the same K_{16} or K_8 blocks. Then $IS_4(8m) = JS_4(8m)$.

The only difference for order $8m + 1$ is that, since $IS_4(9)$ includes all intersection numbers expected, including “ $b - 2$ ”, we may merely use a $\{K_9, K_{4,4}\}$ -decomposition of K_{8m+1} , in order to achieve $IS_4(8m + 1) = JS_4(8m + 1)$.

We have now proved

THEOREM 3.2 *The intersection numbers for S_4 -designs are given by $IS_4(n) = \{0, 1, \dots, b - 2, b\}$ where $n \equiv 0$ or $1 \pmod{8}$, except that $5 \notin IS_4(8)$. \square*

4 D , a triangle with pendant edge

Once again, since D has four edges, we find that a D -design of order n contains $n(n - 1)/8$ blocks and so $n \equiv 0$ or $1 \pmod{8}$. However, since D contains an odd cycle (a triangle!) there is no D -decomposition of any bipartite graph, so in this case we require a D -decomposition of a tripartite graph.

EXAMPLE 4.1 $ID(K_{2,2,2}) \supseteq \{0, 3\}$, and $ID(K_{4,4,4}) \supseteq \{0, 3, 6, 9, 12\}$.

For $K_{2,2,2}$, take the vertex sets $\{1, 1'\} \cup \{2, 2'\} \cup \{3, 3'\}$. Then disjoint D -decompositions are given by $\{(1, 3, 2)-1', (3, 2', 1')-3', (1, 2', 3')-2\}$ and $\{(1', 3', 2')-1, (3', 2, 1)-3, (1', 2, 3)-2'\}$. Thus $\{0, 3\} \subseteq ID(K_{2,2,2})$.

Now let the vertex sets for $K_{4,4,4}$ be $\{A, D\} \cup \{B, E\} \cup \{C, F\}$, where each letter here is itself a set of two points. Then we may take four decompositions of $K_{2,2,2}$ on the four sets $A \cup B \cup F, A \cup E \cup C, D \cup B \cup C$ and $D \cup E \cup F$, yielding 12 blocks for a D -decomposition of $K_{4,4,4}$. Then using the intersection values for $ID(K_{2,2,2})$ we obtain $ID(K_{4,4,4}) \supseteq \{0, 3, 6, 9, 12\}$. \square

For the general construction, we take the vertex set $V = \{(i, j) \mid 1 \leq i \leq 2m, 1 \leq j \leq 4\}$ if $n = 8m$, or $V \cup \{\infty\}$ if $n = 8m + 1$.

Then if $2m \equiv 0$ or 2 (mod 6), $2m \geq 6$, we may use a GDD with group size 2 and block size 3 on $\{1, 2, \dots, 2m\}$, while if $2m \equiv 4$ (mod 6), $2m \geq 10$, we may use a GDD with one group of size 4 and the rest of size 2, and block size 3 on $\{1, 2, \dots, 2m\}$. These exist; see for instance Lemma 2.1 in [1], or the general result in [7]. Then for each group $\{x_1, \dots, x_g\}$ of the GDD, place a D -design on the set $\{(x_i, j) \mid 1 \leq i \leq g, 1 \leq j \leq 4\}$ or on this set together with ∞ . Since the group sizes are 2 or 4, this means we require D -designs of orders 8, 9, 16 and 17. And for each block $\{a, b, c\}$ of the GDD, place a D -decomposition of $K_{4,4,4}$ on $\{(a, j) \mid 1 \leq j \leq 4\} \cup \{(b, j) \mid 1 \leq j \leq 4\} \cup \{(c, j) \mid 1 \leq j \leq 4\}$.

It now remains to deal with orders 8, 9, 16 and 17.

EXAMPLE 4.2 $ID(8) = \{0, 1, \dots, 5, 7\}$.

Take the vertex set $\{\infty\} \cup \mathbb{Z}_7$, and blocks $B = \{(i, 1+i, 3+i)-\infty \mid i \in \mathbb{Z}_7\}$. Note the following trades.

- $X = \{(1, 2, 4)-\infty, (3, 4, 6)-\infty\}$ trades with $X' = \{(1, 2, 4)-3, (\infty, 4, 6)-3\}$,
- $Y = \{(2, 3, 5)-\infty, (4, 5, 0)-\infty\}$ trades with $Y' = \{(2, 3, 5)-4, (\infty, 5, 0)-4\}$,
- $Z = \{(5, 6, 1)-\infty, (0, 1, 3)-\infty\}$ trades with $Z' = \{(5, 6, 1)-0, (\infty, 1, 3)-0\}$,
- $A = \{(0, 1, 3)-\infty, (2, 3, 5)-\infty, (5, 6, 1)-\infty\}$ trades with
- $A' = \{(0, 3, 1)-\infty, (2, 5, 3)-\infty, (6, 1, 5)-\infty\}$.

Here X, Y and Z are pairwise disjoint, and A is also disjoint from X . Thus we achieve the following intersection values, where α below denotes the permutation (1∞) applied to B .

trades	blocks changed	intersection achieved
$B\alpha$	7	0
X, Y, Z	6	1
X, A	5	2
X, Y	4	3
A	3	4
X	2	5
nothing	0	7

\square

EXAMPLE 4.3 $ID(9) = \{0, 1, \dots, 7, 9\}$.

With vertex set \mathbb{Z}_9 , let $D = \{(i, i + 1, i + 4) - (i + 6) \mid i \in \mathbb{Z}_9\}$. The following trades are disjoint:

$$\begin{aligned} X &= \{(1, 2, 5) - 7, (4, 5, 8) - 1\} \text{ trades with } X' = \{(8, 4, 5) - 7, (2, 5, 1) - 8\}, \\ Y &= \{(2, 3, 6) - 8, (5, 6, 0) - 2\} \text{ trades with } Y' = \{(3, 6, 2) - 0, (0, 5, 6) - 8\}, \\ Z &= \{(0, 1, 4) - 6, (3, 4, 7) - 0, (6, 7, 1) - 3\} \text{ trades with} \\ Z' &= \{(3, 1, 7) - 6, (0, 7, 4) - 3, (6, 4, 1) - 0\}. \end{aligned}$$

Now denote permutations by $\alpha = (01)$, $\beta = (125)$, $\gamma = (1234)$, and let $T = \{(7, 8, 2) - 4, (8, 0, 3) - 5\}$. The following table then completes this example.

blocks	intersection size
$D \cap D\gamma$	0
$D \cap D\beta$	1
$D \cap \{X' \cup Y' \cup Z' \cup T\}$	2
$D \cap D\alpha$	3
$D \cap \{X \cup Y' \cup Z' \cup T\}$	4
$D \cap \{X' \cup Y' \cup Z \cup T\}$	5
$D \cap \{X \cup Y \cup Z' \cup T\}$	6
$D \cap \{X' \cup Y \cup Z \cup T\}$	7
$D \cap D$	9

□

EXAMPLE 4.4 $ID(16) = \{0, 1, \dots, 28, 30\}$.

With vertex set $\mathbb{Z}_{15} \cup \{\infty\}$, a design is given by

$$\{(i, 1 + i, 6 + i) - (8 + i), (i, 3 + i, 7 + i) - \infty\} \text{ where } i \in \mathbb{Z}_{15}.$$

Now blocks A_i trade with A'_i for $0 \leq i \leq 6$ where

$$\begin{aligned} A_i &= \{(i, 3 + i, 7 + i) - \infty, (7 + i, 10 + i, 14 + i) - \infty\} \text{ and} \\ A'_i &= \{(i, 3 + i, 7 + i) - (10 + i), (7 + i, \infty + i, 14 + i) - (10 + i)\}. \end{aligned}$$

Disjoint from these trades are the following five trades, B_i with B'_i , for $0 \leq i \leq 4$, where

$$B_i = \{(i, 1 + i, 6 + i) - (8 + i), (5 + i, 6 + i, 11 + i) - (13 + i), (10 + i, 11 + i, 1 + i) - (3 + i)\}$$

and

$$B'_i = \{(i, 6 + i, 1 + i) - (3 + i), (5 + i, 11 + i, 6 + i) - (8 + i), (10 + i, 1 + i, 11 + i) - (13 + i)\}$$

(addition in \mathbb{Z}_{15}). Thus we have trades on $2a + 3b$ blocks, where $0 \leq a \leq 7$ and $0 \leq b \leq 5$. This means that we may trade $2a + 3b = c$ blocks for $2 \leq c \leq 29$. Thus $\{1, 2, \dots, 28\} \subseteq ID(16)$. And trivially $30 \in ID(16)$. Finally, to show $0 \in ID(16)$, let

$$X = \{(6, 9, 13) - \infty, (13, 1, 5) - \infty, (14, 2, 6) - \infty\}$$

which trades with

$$X' = \{(14, 2, 6) - 9\lambda, (1, 15, 13) - 9, (13, 6, \infty) - 15\}.$$

Thus trading $\{B_i\}_{i=0}^4 \cup \{A_i\}_{i=0}^5 \cup \{X\}$ will change *all* the blocks, so $0 \in ID(16)$. This concludes the example. \square

EXAMPLE 4.5 $ID(17) = \{0, 1, \dots, 32, 34\}$.

Let the vertex set be \mathbb{Z}_{17} . Then a design is given by

$$D = \{(i, i + 3, i + 8) - (i + 12), (i, i + 1, i + 7) - (i + 9) \mid i \in \mathbb{Z}_{17}\}.$$

Let permutations on \mathbb{Z}_{17} be given by

$$\begin{aligned} \alpha_0 &= (0\ 1\ 2\ 3\ 4\ 5\ 6\ 7\ 8), & \alpha_1 &= (0\ 1\ 2\ 3\ 4\ 5\ 6\ 7\ 8\ 9\ 10), \\ \alpha_2 &= (0\ 1)(2\ 3\ 4\ 5\ 6), & \alpha_3 &= (0\ 1)(2\ 3\ 4\ 5), & \alpha_4 &= (0\ 1\ 2\ 3\ 4). \end{aligned}$$

Then $|D \cap D\alpha_i| = i$, $0 \leq i \leq 4$, so $\{0, 1, 2, 3, 4\} \subseteq ID(17)$. For the remaining intersection values we consider trades as follows.

The set $A_i = \{(1, 4, 9) - 13, (13, 16, 4) - 8\} + i \pmod{17}$ trades with $A'_i = \{(16, 4, 13) - 9, (9, 1, 4) - 8\} + i \pmod{17}$, $0 \leq i \leq 4$. Disjoint from this are the blocks

$$B_i = \{(1, 2, 8) - 10, (9, 10, 16) - 1\} + i \pmod{17}$$

trading with

$$B'_i = \{(8, 2, 1) - 16, (9, 16, 10) - 8\} + i \pmod{17},$$

$0 \leq i \leq 7$. Also let

$$C_i = \{(0, 3, 8) - 12, (12, 15, 3) - 7, (11, 12, 1) - 3\} + i,$$

which trades with

$$C'_i = \{(0, 8, 3) - 7, (12, 15, 3) - 1, (1, 11, 12) - 8\} + i,$$

for $0 \leq i \leq 4$.

Note that C_0 is disjoint from A_i , $i = 0, 1, 2, 3$,
 C_1 is disjoint from A_i , $i = 1, 2, 3, 4$,
 C_2 is disjoint from A_i , $i = 0, 2, 3, 4$,
 C_3 is disjoint from A_i , $i = 0, 1, 3, 4$,
 C_4 is disjoint from A_i , $i = 0, 1, 2, 4$.

Thus we may obtain trades of sizes 2, 3, ..., 28, 29, yielding $\{5, 6, \dots, 31, 32\} \subseteq ID(17)$. Finally, $34 \in ID(17)$ trivially. This completes the example. \square

Now combining the results of this section we have

THEOREM 4.1 *The intersection numbers for D-designs are given by $ID(n) = \{0, 1, \dots, b - 2, b\}$ where $b = n(n - 1)/8$.* \square

5 The graph Y

A Y -design of order n contains $n(n-1)/8$ blocks, and so $n \equiv 0$ or $1 \pmod{8}$. The only ingredients we need are Y -designs of orders 8 and 9, a Y -decomposition of $K_{4,4}$, and their intersection numbers. (In fact, it suffices to use $IY(K_{4,4}) \supseteq \{0, 4\}$.)

EXAMPLE 5.1 $IY(K_{4,4}) \supseteq \{0, 4\}$.

Let the vertex set be $\{1, 2, 3, 4\} \cup \{5, 6, 7, 8\}$. Then two disjoint decompositions are given by

$$\{(4, 7, 1; 5, 6), (1, 8, 2; 6, 7), (2, 5, 3; 7, 8), (3, 6, 4; 5, 8)\}$$

and

$$\{(8, 3, 5; 1, 2), (5, 4, 6; 2, 3), (6, 1, 7; 3, 4), (7, 2, 8; 1, 4)\}.$$

□

EXAMPLE 5.2 $IY(8) = \{0, 1, 2, 3, 4, 5, 7\}$.

With vertex set $\{\infty\} \cup \mathbb{Z}_7$, take blocks $D = A \cup B \cup C$ where

$$A = \{(0, 1, 3; 6, \infty), (1, 2, 4; 0, \infty)\}, \quad B = \{(2, 3, 5; 1, \infty), (3, 4, 6; 2, \infty)\},$$

$$C = \{(4, 5, 0; 3, \infty), (5, 6, 1; 4, \infty), (6, 0, 2; 5, \infty)\}.$$

Blocks A trade with $A' = \{(6, 3, 1; 0, 2), (3, \infty, 4; 0, 2)\}$,

blocks B trade with $B' = \{(1, 5, 3; 2, 4), (5, \infty, 6; 2, 4)\}$ and

blocks C trade with $C' = \{(4, 5, 0; 3, 2), (4, 1, 6; 5, 0), (5, 2, \infty; 1, 0)\}$.

Now let α denote the permutation (01) and β the permutation (012) . We obtain the following intersection numbers, which completes the result.

blocks	intersection
$D \cap D\beta$	0
$D \cap D\alpha$	1
$D \cap \{A \cup B' \cup C'\}$	2
$D \cap \{A' \cup B' \cup C\}$	3
$D \cap \{A \cup B \cup C'\}$	4
$D \cap \{A \cup B' \cup C\}$	5
$D \cap D$	7

□

EXAMPLE 5.3 $IY(9) = \{0, 1, \dots, 7, 9\}$.

Let the vertex set be \mathbb{Z}_9 , and blocks be $D = \{(0+i, 1+i, 3+i, 6+i, 7+i) \mid i \in \mathbb{Z}_9\}$ (addition mod 9). The blocks

$$A_i = \{(i-1, i, 2+i, 5+i, 6+i), (i, 1+i, 3+i, 6+i, 7+i)\}, \quad 1 \leq i \leq 8,$$

trade with

$$A'_i = \{(5 + i, 2 + i, i; i - 1, i + 1), (2 + i, 6 + i, 3 + i; 1 + i, 7 + i)\}, 1 \leq i \leq 4.$$

Also the blocks

$$B_i = \{(3i - 3, 3i - 2, 3i; 3i + 3, 3i + 4), \\ (3i - 2, 3i - 1, 3i + 1; 3i + 4, 3i + 5), (3i - 1, 3i, 3i + 2; 3i + 5, 3i + 6)\},$$

$1 \leq i \leq 3$, trade with the blocks

$$B'_i = \{(3i + 3, 3i, 3i - 2; 3i - 3, 3i - 1), \\ (3i, 3i + 4, 3i + 1; 3i - 1, 3i + 5), (3i + 2, 3i + 5, 3i + 1; 3i + 4, 3i - 1)\},$$

$1 \leq i \leq 3$. Thus we obtain the required intersection numbers:

blocks	intersection
$D \cap \{B'_1 \cup B'_2 \cup B'_3\}$	0
$D \cap \{((8, 0, 2; 5, 6)) \cup \{A'_i \mid 1 \leq i \leq 4\}\}$	1
$D \cap \{A_1 \cup A'_2 \cup A'_3 \cup B'_3\}$	2
$D \cap \{A'_1 \cup A'_2 \cup A'_3 \cup B_3\}$	3
$D \cap \{A_1 \cup A_2 \cup A'_3 \cup B'_3\}$	4
$D \cap \{A'_1 \cup A'_2 \cup A_3 \cup B_3\}$	5
$D \cap \{A_1 \cup A_2 \cup A_3 \cup B'_3\}$	6
$D \cap \{A'_1 \cup A_2 \cup A_3 \cup B_3\}$	7
$D \cap D$	9

□













Thanks to Lemma 1.1 we now have

THEOREM 5.1 *The intersection numbers for Y -designs are given by $IY(n) = \{0, 1, \dots, b - 2, b\}$ where $b = n(n - 1)/8$.* □

6 Summary

The following table summarises the intersection results for G -designs where G is a connected graph on at most four vertices or at most four edges.

In the table, b denotes the number of blocks in a G -design of order n , and the impossible intersection values are $b - x$ where x is as given. A reference is listed if the result is not in this paper.

G	b	x	Comments	Ref
K_2 	$n(n-1)/2$	all except b	unique design!	
P_3 	$n(n-1)/4$	1	$n \equiv 0, 1 \pmod{4}$	
P_4 	$n(n-1)/6$	1	$n \equiv 0, 1 \pmod{3}$, $n \geq 4$	
P_5 	$n(n-1)/8$	1	$n \equiv 0, 1 \pmod{8}$	
K_3 	$n(n-1)/6$	1, 2, 3, 5	$n \equiv 1, 3 \pmod{6}$, $5, 8 \notin IK_3(9)$.	[8]
D 	$n(n-1)/8$	1	$n \equiv 0, 1 \pmod{8}$	
Y 	$n(n-1)/8$	1	$n \equiv 0, 1 \pmod{8}$	
S_3 	$n(n-1)/6$	1	$n \geq 6$, $n \equiv 0, 1 \pmod{8}$, $3 \notin IS_3(6)$	
S_4 	$n(n-1)/8$	1	$n \equiv 0, 1 \pmod{8}$, $5 \notin IS_4(8)$	
C_4 	$n(n-1)/8$	1	$n \equiv 1 \pmod{8}$	[4]
$K_4 - e$ 	$n(n-1)/10$	1, 2	$n \equiv 0, 1 \pmod{5}$, $n \geq 6$; $7, 8 \notin I(11)$	[5]
K_4 	$n(n-1)/12$	1, 2, 3, 4, 5, 7	$n \equiv 1, 4 \pmod{12}$; $7, 9, 10, 11, 14 \notin I(16)$; several unknown values for $n = 25, 28, 37$.	[6]

References

- [1] P. Adams, E.J. Billington and C.C. Lindner, *k*-perfect $3k$ -cycle systems, J. Combin. Math. Combin. Computing **15** (1994), 141–154.
- [2] J.C. Bermond and J. Schönheim, *G*-decomposition of K_n , where *G* has four vertices or less, Discrete Math. **19** (1977), 113–120.
- [3] E.J. Billington, *The intersection problem for combinatorial designs*, Congressus Numerantium **92** (1993), 33–54.
- [4] E.J. Billington, *The intersection problem for m -cycle systems*, J. Combinatorial Designs **1** (1993), 435–452.
- [5] E.J. Billington, M. Gionfriddo and C.C. Lindner, *The intersection problem for $K_4 - e$ designs*, J. Stat. Plann. Inference (to appear).
- [6] C.J. Colbourn, D.G. Hoffman and C.C. Lindner, *Intersections of $S(2, 4, v)$ designs*, Ars Combinatoria **33** (1992), 97–111.
- [7] C.J. Colbourn, D.G. Hoffman and R. Rees, *A new class of group divisible designs with block size three*, J. Combinatorial Theory, series A **59** (1992), 73–89.
- [8] C.C. Lindner and A. Rosa, *Steiner triple systems having a prescribed number of triples in common*, Canad. J. Math. **27** (1975), 1166–1175. Corrigendum: Canad. J. Math. **30** (1978), 896.

(Received 15/2/95)