

The structure of strong tournaments containing exactly one out-arc pancyclic vertex*

QIAOPING GUO[†] GAOKUI XU

*School of Mathematical Sciences
Shanxi University
Taiyuan, 030006
China*

Abstract

An arc in a tournament T with $n \geq 3$ vertices is called pancyclic if it belongs to a cycle of length l for all $3 \leq l \leq n$. We call a vertex u of T an out-arc pancyclic vertex of T if each out-arc of u is pancyclic in T . Yao, Guo and Zhang [*Discrete Appl. Math.* 99 (2000), 245–249] proved that every strong tournament contains at least one out-arc pancyclic vertex, and they gave an infinite class of strong tournaments, each of which contains exactly one out-arc pancyclic vertex. In this paper we give the structure of strong tournaments containing exactly one out-arc pancyclic vertex.

1 Introduction

Let D be a digraph with the vertex set $V(D)$ and the arc set $A(D)$. We denote the number of vertices in D by $|V(D)|$. A subdigraph induced by a subset $A \subseteq V(D)$ is denoted by $D[A]$. We also write $D - A$ for $D[V(D) - A]$.

A *tournament* is a digraph, where there is precisely one arc between every pair of distinct vertices. An *l -cycle* is a cycle of length l . An arc in a digraph D is said to be *pancyclic*, if it belongs to an l -cycle for all $3 \leq l \leq n$. An arc leaving from a vertex x in a digraph is called an *out-arc* of x . We call a vertex u of T an out-arc pancyclic vertex of T , if each out-arc of u is pancyclic in T .

A digraph D is said to be *strong*, if for every pair of vertices x and y , D contains a path from x to y and a path from y to x . A directed path from x to y in D is

* This work is supported by the Natural Science Young Foundation of China (Nos.11201273, 11401352, 11401354) and by the Natural Science Foundation of Shanxi Province, China (No.2016011005).

[†] Corresponding author, guoqp@sxu.edu.cn.

denoted by (x, y) -path. D is called k -strong if $|V(D)| \geq k + 1$ and $D - X$ is strong for any set $X \subseteq V(D)$ with $|X| < k$. If a digraph is strong, it is 1-strong. If a digraph D is k -strong, but not $(k + 1)$ -strong, then $\kappa(D) = k$ is defined as the *strong connectivity* of D .

In [4], Thomassen confirmed that every strong tournament contains a vertex x such that each out-arc of x is contained in a Hamilton cycle. In 2000, Yao et al.[5] extended the result of Thomassen and proved that every strong tournament T has an out-arc pancyclic vertex. In the same paper, they also found an infinite class of strong tournaments, each of which contains exactly one out-arc pancyclic vertex (see Figure 1). For a strong tournament T with minimum out-degree at least two, Guo et al. [3] proved that T contains at least three out-arc pancyclic vertices. This implies that a strong tournament containing exactly one out-arc pancyclic vertex can only be 1-strong.

Example. (Yao [5]). Let $n \geq 5$ be an integer and let T_n be a tournament with the vertex set $\{v_1, v_2, \dots, v_n\}$ and the arc set

$$\{v_i v_j \mid 2 \leq i < j \leq n\} \cup \{v_{n-1} v_1, v_n v_1\} \cup \{v_1 v_j \mid 2 \leq j \leq n - 2\}.$$

Then T_n contains exactly one vertex v_n whose out-arcs are pancyclic.




Figure 1: A strong tournament T containing exactly one out-arc pancyclic vertex, where $v_i \rightarrow v_j$ for all $2 \leq i < j \leq n$.

In this paper, we obtain a sufficient and necessary condition for a strong tournament to contain exactly one out-arc pancyclic vertex.

2 Terminology and Preliminaries

We assume that the reader is familiar with the standard terminology on digraphs and refer the reader to [1].

If uv is an arc of a digraph D , then we say that u dominates v or uv is an *out-arc* of u , and we write $u \rightarrow v$. For disjoint subsets X and Y of $V(D)$, if every vertex of X dominates every vertex of Y , we say X dominates Y and write $X \rightarrow Y$. If there is no arc from a vertex in Y to a vertex in X , we say $X \Rightarrow Y$. For a tournament, $X \rightarrow Y$ if and only if $X \Rightarrow Y$.

Let $x \in V(D)$ and H be a subdigraph of D . The set of all vertices in H dominating x (dominated by x , respectively) is denoted by $N_H^-(x)$ ($N_H^+(x)$, respectively). Furthermore, $d_H^-(x) = |N_H^-(x)|$ ($d_H^+(x) = |N_H^+(x)|$, respectively) is called the *in-degree* (*out-degree*, respectively) of x in H . We will omit the subscript if $H = D$. We use $\delta^+(D) = \min\{d_D^+(x) \mid x \in V(D)\}$ to stand for the *minimum out-degree* of D .

A path (cycle) containing every vertex of D is called a *Hamilton path* (*Hamilton cycle*) of D . For two vertices u, v on a path P , we let uPv denote the unique (u, v) -path on P .

A set $X \subseteq V(D)$ is called a *separating set* if $D - X$ is not strong. A vertex $u \in V(D)$ is also called a *cut vertex* of D if $D - \{u\}$ is not strong. A *strong component* of a digraph D is a maximal induced subdigraph of D which is strong. If D is not strong, then we can partition the vertices in D into strong components T_1, T_2, \dots, T_r ($r \geq 2$), such that $T_i \Rightarrow T_j$ if and only if $i < j$.

In the proofs of our main results, the following results are very useful.

Lemma 2.1 (Camion [2]). *A non-trivial tournament is strong if and only if it has a Hamilton cycle.*

Lemma 2.2 (Yao [5]). *Let T_n be a strong tournament on n vertices and assume that the vertices of T_n are labeled u_1, u_2, \dots, u_n such that $d^+(u_1) \leq d^+(u_2) \leq \dots \leq d^+(u_n)$. If $d^+(u_1) = 1$, then all out-arcs of u_1 are pancyclic.*

Lemma 2.3 (Guo [3]). *Every strong tournament T with minimum out-degree at least two contains at least three out-arc pancyclic vertices.*

Lemma 2.4 (Yeo [6]). *Let D be a k -strong digraph, with $k \geq 1$, and let S be a separating set in D , such that $T = D - S$ is a tournament. Let T_1, T_2, \dots, T_r ($r \geq 2$) be the strong components of T , such that $T_1 \Rightarrow T_2 \Rightarrow \dots \Rightarrow T_r$. Then, for every $1 \leq l \leq |V(T)| - 1$, $u \in T_1$ and $v \in T_r$, there exists a (u, v) -path of length l in T .*

3 Main results

Theorem 3.1. *Let T be a strong tournament with strong connectivity $\kappa(T) = 1$. Let v be a cut vertex of T and let T_1, T_2, \dots, T_r ($r \geq 2$) be the strong components of $T - v$, such that $T_i \Rightarrow T_j$ if and only if $i < j$. Let $x \in V(T_{r-1})$. Then the following hold:*

- (1) *If $V(T_{r-1}) = \{x\}$, then x is an out-arc pancyclic vertex of T if and only if $V(T_r) \rightarrow v \rightarrow x$.*
- (2) *If $|V(T_{r-1})| > 1$, then x is an out-arc pancyclic vertex of T if and only if $V(T_r) \rightarrow v \rightarrow x$ and for each $y \in N_{T_{r-1}}^+(x)$, there exists a Hamilton path containing xy in T_{r-1} and there exists a 3-cycle containing xy in $D[V(T_{r-1}) \cup \{v\}]$.*

Proof. Since T is strong, there is a vertex u in T_1 such that $v \rightarrow u$.

(1) (Sufficiency.) Suppose $V(T_{r-1}) = \{x\}$ and $V(T_r) \rightarrow v \rightarrow x$. Let $x_1 \in N^+(x)$ be arbitrary. Then $x_1 \in V(T_r)$ and $x_1 \rightarrow v$. It is clear that xx_1vx is a 3-cycle containing xx_1 . Using Lemma 2.1 on each component, it is easy to prove that there exists a (u, x) -path P_l of length l in T for every $1 \leq l \leq |V(T)| - |V(T_r) \cup \{v\}| - 1$. Thus $xx_1v u P_l x$ is a cycle of length $l + 3$ containing xx_1 for every $1 \leq l \leq |V(T)| - |V(T_r)| - 2$. Suppose that $x_1x_2 \dots x_{|V(T_r)|}x_1$ is a Hamilton cycle of T_r . Then $xx_1x_2 \dots x_i v u P_{|V(T)| - |V(T_r)| - 2} x$ is a cycle of length $|V(T)| - |V(T_r)| + i$ containing xx_1 for every $2 \leq i \leq |V(T_r)|$. Therefore, xx_1 is contained in an m -cycle for every $m \in \{3, 4, \dots, |V(T)|\}$, we note that xx_1 is pancyclic and x is an out-arc pancyclic vertex of T .

(Necessity.) Now, let $V(T_{r-1}) = \{x\}$ and x is an out-arc pancyclic vertex of T . We will prove $V(T_r) \rightarrow v \rightarrow x$. Let $z \in V(T_r)$ be arbitrary. Then $z \in N^+(x)$ and xz is pancyclic in T . So xz is in a 3-cycle. Note that $T_i \Rightarrow T_j$ if and only if $i < j$. This implies that $z \rightarrow v \rightarrow x$. That is, we have that $V(T_r) \rightarrow v \rightarrow x$.

(2) (Sufficiency.) Suppose that $|V(T_{r-1})| > 1$, $V(T_r) \rightarrow v \rightarrow x$ and for each $y \in N^+_{T_{r-1}}(x)$, there exists a Hamilton path containing xy in T_{r-1} and there exists a 3-cycle containing xy in $D[V(T_{r-1}) \cup \{v\}]$. We will prove that x is an out-arc pancyclic vertex of T . Let $w \in N^+(x)$ be arbitrary. Then $w \in V(T_{r-1}) \cup V(T_r)$. We only need to prove that xw is pancyclic in T .

If $w \in V(T_{r-1})$, by assumption, there exists a Hamilton path containing xw in T_{r-1} and there exists a 3-cycle containing xw in $D[V(T_{r-1}) \cup \{v\}]$. Using Lemma 2.1 on each component except T_{r-1} and using the Hamilton path containing xw in T_{r-1} , it is easy to prove that, for each $z \in V(T_r)$, there exists a (u, z) -path P_l of length l in T containing xw for every $3 \leq l \leq |V(T)| - 2$. Thus $u P_l z v u$ is a cycle of length $l + 2$ containing xw for every $3 \leq l \leq |V(T)| - 2$. Note that $xwzvx$ is a 4-cycle containing xw for any $z \in V(T_r)$ and there exists a 3-cycle containing xw in $D[V(T_{r-1}) \cup \{v\}]$. Therefore, xw is contained in an m -cycle for every $m \in \{3, 4, \dots, |V(T)|\}$, we note that xw is pancyclic.

If $w \in V(T_r)$, then $w \rightarrow v$. It is clear that $xwvx$ is a 3-cycle containing xw . Using Lemma 2.1 on each component except T_r , it is easy to prove that there exists a (u, x) -path P_l of length l in T for every $1 \leq l \leq |V(T)| - |V(T_r) \cup \{v\}| - 1$. Thus $xwv u P_l x$ is a cycle of length $l + 3$ containing xw for every $1 \leq l \leq |V(T)| - |V(T_r)| - 2$. Suppose that $x_1x_2 \dots x_{|V(T_r)|}$ is a Hamilton cycle of T_r with $x_1 = w$. Then $xx_1x_2 \dots x_i v u P_{|V(T)| - |V(T_r)| - 2} x$ is a cycle of length $|V(T)| - |V(T_r)| + i$ containing xw for every $2 \leq i \leq |V(T_r)|$. Therefore xw is contained in an m -cycle for every $m \in \{3, 4, \dots, |V(T)|\}$, and we note that xw is pancyclic.

(Necessity.) Let $x \in V(T_{r-1})$ be an out-arc pancyclic vertex of T . We first prove $V(T_r) \rightarrow v \rightarrow x$. Let $z \in V(T_r)$ be arbitrary. Then $z \in N^+(x)$ and xz is pancyclic in T . So xz is in a 3-cycle which implies that $z \rightarrow v \rightarrow x$. That is, we have that $V(T_r) \rightarrow v \rightarrow x$.

For each $y \in N^+_{T_{r-1}}(x)$, we have that xy is pancyclic in T , and so xy is contained in an m -cycle for every $m \in \{3, 4, \dots, |V(T)|\}$. Let $C = vx_1x_2 \dots x_{|V(T)|-1}v$ be a

Hamilton cycle containing xy with some $x_k = x$ and $x_{k+1} = y$. Suppose that starting from v , x_i is the first vertex and x_j is the last one appearing in $V(T_{r-1})$ on C . Then $i \leq k < j$ and $x_i x_{i+1} \dots x_j$ is a Hamilton path containing xy of T_{r-1} . Since xy is in a 3-cycle, it is clear that such a 3-cycle can only be contained in the $D[V(T_{r-1}) \cup \{v\}]$. \square

Theorem 3.2. *Let T be a strong tournament with n vertices and strong connectivity $\kappa(T)$. Then T contains exactly one out-arc pancyclic vertex if and only if all of the following hold.*

- (1) $n \geq 5$.
- (2) $\kappa(T) = 1$ and there exists a cut vertex v of T and strong components T_1, T_2, \dots, T_r ($r \geq 3$) of $T - v$ such that $T_i \Rightarrow T_j$ if and only if $i < j$ and $|V(T_r)| = 1$.
- (3) There exists a vertex $z \in V(T_2) \cup \dots \cup V(T_{r-1})$ such that $v \rightarrow z$.
- (4) If $V(T_{r-1}) = \{u_1\}$, then $u_1 \rightarrow v$. If $|V(T_{r-1})| > 1$, then, for any $x \in V(T_{r-1})$, we have $x \rightarrow v$ or there exists a vertex $y \in N_{T_{r-1}}^+(x)$ such that there is no Hamilton path containing xy in T_{r-1} or there exists a vertex $z \in N_{T_{r-1}}^+(x)$ such that there is no 3-cycle containing xz in $D[V(T_{r-1}) \cup \{v\}]$.

Proof. (Sufficiency.) Let T be a strong tournament with n vertices and strong connectivity $\kappa(T)$ and T satisfies (1)–(4). By (2), assume that v is a cut vertex of T and T_1, T_2, \dots, T_r ($r \geq 3$) are strong components of $T - v$ such that $T_i \Rightarrow T_j$ if and only if $i < j$ and $|V(T_r)| = 1$. Let $V(T_r) = \{u\}$. Then $d^+(u) = 1$ and Lemma 2.2 implies that u is an out-arc pancyclic vertex of T . We will prove that any vertex except u in T is not an out-arc pancyclic vertex of T . By (3), there exists a vertex $z \in V(T_2) \cup \dots \cup V(T_{r-1})$ such that $v \rightarrow z$. Then vz can not be contained in any Hamilton cycle of T and so v is not an out-arc pancyclic vertex of T . Let $x \in V(T_1) \cup \dots \cup V(T_{r-2})$ be arbitrary. It is easy to see that xu can not be contained in any Hamilton cycle of T and so x is also not an out-arc pancyclic vertex of T . We consider the vertices of T_{r-1} . By (4) and Theorem 3.1, we have that any vertex T_{r-1} is not an out-arc pancyclic vertex of T . Thus, u is an unique out-arc pancyclic vertex of T .

(Necessity.) Let T be a strong tournament with n vertices and strong connectivity $\kappa(T)$ which contains exactly one out-arc pancyclic vertex. By lemma 2.3, T is a strong tournament with minimum out-degree 1. Let M be the set of vertices with out-degree 1 in T . By Lemma 2.2, all vertices in M are out-arc pancyclic vertices of T . So we have $|M| = 1$. If $n \leq 4$, then it is clear that $|M| \geq 2$, a contradiction. Therefore, we have $n \geq 5$ and (1) is proved.

Let $M = \{u\}$ and $N^+(u) = \{v\}$. Then v is a cut vertex of T and $\kappa(T) = 1$. Let T_1, T_2, \dots, T_r ($r \geq 2$) be the strong components of $T - v$, such that $T_i \Rightarrow T_j$ if and only if $i < j$. Obviously, $V(T_r) = \{u\}$ and u is an out-arc pancyclic vertex of T . If $r = 2$, it is easy to see that v is also an out-arc pancyclic vertex of T , a contradiction. So we have $r \geq 3$. (2) is proved.

If $V(T_2) \cup \cdots \cup V(T_{r-1}) \Rightarrow \{v\}$, then $N^+(v) \subseteq V(T_1)$. Let $v_1 \in N^+(v)$ be arbitrary. By Lemma 2.4, there exists a (v_1, u) -path P_l of length l in T for every $1 \leq l \leq |V(T)| - 2$. So vv_1P_luv is an $(l + 2)$ -cycle containing vv_1 for every $1 \leq l \leq |V(T)| - 2$, which implies that v is an out-arc pancyclic vertex of T , a contradiction. Therefore, there exists a vertex $z \in V(T_2) \cup \cdots \cup V(T_{r-1})$ such that $v \rightarrow z$ and (3) is proved.

Since u is the unique out-arc pancyclic vertex of T , we have that no vertex in T_{r-1} is an out-arc pancyclic vertex of T . Then (4) is obvious by using Theorem 3.1. \square

References

- [1] J. Bang-Jensen and G. Gutin, *Digraphs: Theory, Algorithms and Applications*, Springer, London, 2000.
- [2] P. Camion, Chemins et circuits hamiltoniens des graphes complets, *C. R. Acad. Sci. Paris* 249 (1959), 2151–2152.
- [3] Q. Guo, S. Li, H. Li and H. Zhao, The numbers of vertices whose out-arcs are pancyclic in a strong tournament, *Graphs Combin.* 30 (2014), 1163–1173.
- [4] C. Thomassen, Hamiltonian-connected tournaments, *J. Combin. Theory Ser. B* 28 (1980), 142–163.
- [5] T. Yao, Y. Guo and K. Zhang, Pancyclic out-arcs of a vertex in a tournament, *Discrete Appl. Math.* 99 (2000), 245–249.
- [6] A. Yeo, The number of pancyclic arcs in a k -strong tournament, *J. Graph Theory* 50 (2005), 212–219.

(Received 16 Nov 2016)